

Egg viability decreases rapidly with time since ovulation in the rainbow darter *Etheostoma caeruleum*: implications for the costs of choosiness

R. L. MORAN*, R. M. SOUKUP, M. ZHOU AND R. C. FULLER

Program in Ecology, Evolution, and Conservation Biology, Department of Animal Biology,
School of Integrative Biology, University of Illinois at Urbana-Champaign, Champaign,
Illinois 61820, U.S.A.

(Received 4 October 2017, Accepted 18 November 2017)

Egg viability in the rainbow darter *Etheostoma caeruleum*, a fish apparently lacking female mate choice, was found to decline rapidly after ovulation. It was observed that the majority of a female's clutch may fail to hatch if she is prevented from mating for as little as 6 h. These data suggest that exercising female mate preferences may be selectively disfavoured in *E. caeruleum* due to the high cost of delaying mating.

© 2017 The Fisheries Society of the British Isles

Key words: egg overripening; *Etheostoma caeruleum*; female mate choice; mate choice cost; rainbow darter.

The degree of discrimination exhibited by female animals when choosing a mate ranges from none to extreme. This diversity exists because female mate choice is shaped by a number of variables, including the subset of males made available to her by environmental factors and male–male interactions (Beehler & Foster, 1988; Jennions & Petrie, 1997; Wong & Candolin, 2005) and the costs of mate choice behaviour (Janetos, 1980; Real, 1990; Kokko *et al.*, 2003). If the costs of choosing outweigh the benefits, the most favoured strategy would be to mate with the first male the female encounters, *i.e.* random mating. Several costs to female mate choice (*e.g.* time and energy expenditure, predator exposure) have been investigated across a variety of taxa (Reynolds & Gross, 1990; Jennions & Petrie, 1997), but decline in gametic quality over time has thus far received little attention. In fishes, the phenomenon of decreasing egg viability post-ovulation, termed egg overripening, has been reported across a range of species including Atlantic salmon *Salmo salar* L. 1758 (de Gaudemar & Beall, 1998), goldfish *Carassius auratus* (L. 1758) (Formacion *et al.*, 1993), Atlantic halibut *Hippoglossus hippoglossus* (L. 1758) (Bromage *et al.*, 1994) and turbot *Scophthalmus maximus* (L. 1758) (McEvoy, 1984). Having eggs susceptible to overripening would

*Author to whom correspondence should be addressed. Tel.: +1 217 333 2235; email: rmoran9@illinois.edu.

pressure females to spawn quickly, as waiting could incur a substantial fitness cost. Such selection for rapid mate acquisition may consequently lead to a decrease in female choosiness and influence the expression of female mating preferences.

The rainbow darter *Etheostoma caeruleum* Storer 1845 is a small benthic fish common in freshwater streams across the eastern U.S.A. (Page, 1983). During the breeding season from late March to early June, vividly coloured males attempt to guard gravid females from rival males. Females signal their readiness to spawn by performing nose digs, wherein she pushes her head into the gravel. Spawning involves the female burying herself shallowly in gravel with an arched posture; once a male takes position above her, both vibrate and release eggs and sperm. No parental care is practised (Winn, 1958; Fuller, 2003).

Although there is evidence that female *E. caeruleum* favour some types of males in dichotomous trials, such preferences have little apparent effect on the outcome of mating due to male–male competition (Fuller, 2003). In any case, female *E. caeruleum* demonstrate no overt choice when allowed to interact freely with males; having assumed the spawning position, the female always spawns with the first male to arrive regardless of its characteristics (R. C. Fuller & M. Zhou, pers. obs.), suggesting that exercising choice may be disadvantageous. Good reasons exist to suspect that mate choice in female *E. caeruleum* may harbour a cost in prolonging the time between ovulation and spawning. In previous experiments, females have been observed expelling and subsequently eating unfertilized eggs when held in isolation (Zhou & Fuller, 2014). Furthermore, isolated females subsequently allowed to spawn with males often produce entirely nonviable clutches (R. L. Moran, pers. obs.). This study aimed to formally test the hypothesis that delaying spawning is costly for female *E. caeruleum*, by quantifying change in egg viability as a function of time since ovulation.

Etheostoma caeruleum were collected by kick seine from Mill Pond Outlet (Kalamazoo Co., MI, U.S.A.; c. 42° 20' 13" N; 85° 24' 10" W) in April and May 1998 (year 1) and from an unnamed tributary of the Saline Branch Drainage Ditch (Champaign Co., IL, U.S.A.; c. 40° 07' 40" N; 88° 10' 28" W) in March 2017 (year 2). Fish from year 1 were maintained at the Kellogg Biological Station. Fish from year 2 were maintained at the University of Illinois at Urbana-Champaign. In both years, fish were housed in male–female pairs in 381 aquaria with gravel substratum, maintained at external ambient temperature and light:dark cycle. The fish were fed frozen bloodworms (chironomid larvae) and live tubifex worms twice per day.

Fish were monitored during daylight hours over 4–5 days post-capture. A female was assumed to have recently ovulated when she performed a nose dig. After a female performed a nose dig, she was moved to an empty tank. Females were kept in isolation for various lengths of time: 0 h ($n = 10$), 6 h ($n = 6$), 12 h ($n = 7$), 24 h ($n = 5$). Following the isolation period, a male was introduced. Unique males were used for each female. All males were clearly in breeding condition as shown by their nuptial coloration. Males were visually size matched and care was taken to avoid smaller males that might have low sperm production. After the introduction of the male, the pair were left together for 4 h or until the female finished spawning her clutch. The resulting eggs were collected with a siphon and placed in small tubs filled with water; dilute methylene blue was added to inhibit fungal growth. Hatching success was recorded as the number of eggs yielding fry out of total number of eggs collected. A quasibinomial regression with a logit link function was used to test for a relationship between hatching success




FIG. 1. Mean \pm S.E. proportion of eggs that hatched from a clutch of *Etheostoma caeruleum* eggs decreased with increasing time that a female was held in isolation (i.e. prevented from spawning) after ovulation occurred.

and female isolation time. To determine whether there was an effect of collection year and location on hatching success, year was included as a covariate in the model. The quasibinomial error distribution was used to account for overdispersion in the response variable (i.e. hatching success). Statistical analysis was performed in R 3.4.0 (www.r-project.org).

All females spawned following the introduction of a male. The number of eggs collected ranged from 17 to 110 (Table S1, Supporting information) and was uncorrelated with isolation time or year. Hatching success declined strongly as a function of increasing female isolation time ($F_{1,25} = 5.91, P < 0.05$; Fig. 1). There was no effect of year on hatching success ($F_{1,25} = 3.30, P > 0.05$). Although hatching success varied at each female holding time (Fig. 1), these data suggest that on average, greater than 50% of a female's clutch is likely to become non-viable if retained for as little as 6 h after ovulation.

Given that hatching success declines precipitously over time, gravid female *E. caeruleum* appear to be under strong pressure to fully spawn a clutch of eggs in >24 h. Females that fail to do so risk substantial egg mortality. The time constraint for spawning may be exacerbated by the fact that female *E. caeruleum* release only a small fraction of ovulated eggs per spawning bout and thus must spawn multiple times to fully expel an entire clutch (Fuller, 1998). Under these conditions, the cost for a female *E. caeruleum* to reject a male may be unacceptably high.

Female mate choice in *E. caeruleum* may be further disfavoured by strong male–male competition. Male *E. caeruleum* fight vigorously for access to gravid females, attempting to monopolize spawning and prevent the participation of sneaker males (Winn, 1958; Fuller, 2003). Hence, the choice of males within a single patch is probably limited to those that are competitively superior. Furthermore, there may be

little additional benefit for the female to choose following male–male competition if male competitive ability predicts fitness benefits to females and her offspring (Wong & Candolin, 2005).

Darters are a highly speciose clade that have received increasing attention from evolutionary biologists over the past decade. Spectacular and diverse male colouration in darters has been suggested to act as an agent of speciation by sexual selection, with the most commonly posited mechanism being divergent female mate choice (Mendelson, 2003; Williams & Mendelson, 2010; Williams *et al.*, 2013). Evidence is mounting, however, that female choice is limited in at least some darter species (Pyron, 1995; Fuller, 2003; Moran *et al.*, 2013, 2017; Zhou *et al.*, 2015). This study suggests that female preference in darters is probably costly due to the need to spawn shortly after ovulation while egg viability remains high. Considering that egg overripening seems to be common across a variety of fish species, often occurring over time frames comparable with or shorter than in *E. caeruleum* (Kjørsvik *et al.*, 1990), its role in shaping female mate choice may be underappreciated.

This work was supported by the Cooperative State Research, Education and Extension Service, U.S. Department of Agriculture, under project number ILLU 875-952, the National Science Foundation (DEB 0953716 and IOS 1701676) and the University of Illinois. The treatment of animals was approved by the Institutional Animal Care and Use Committee under protocol #17031.

Supporting Information

Supporting Information may be found in the online version of this paper:

TABLE S1. Isolation time (time held after ovulation), total number of eggs collected, and total number of eggs hatched for each female *Etheostoma caeruleum*.

References

Beehler, B. M. & Foster, M. S. (1988). Hotshots, hotspots and female preference in the organization of lek mating systems. *The American Naturalist* **131**, 203–219. <https://doi.org/10.1086/284786>

Bromage, N., Bruce, M., Basavaraja, N., Rana, K., Shields, R., Young, C., Dye, J., Smith, P., Gillespie, M. & Gamble, J. (1994). Egg quality determinants in finfish: the role of overripening with special reference to the timing of stripping in the Atlantic halibut *Hippoglossus hippoglossus*. *Journal of the World Aquaculture Society* **25**, 13–21. <https://doi.org/10.1111/j.1749-7345.1994.tb00799.x>

Formacion, M. J., Hori, R. & Lam, T. J. (1993). Overripening of ovulated eggs in goldfish: I Morphological changes. *Aquaculture* **114**, 155–168. [https://doi.org/10.1016/0044-8486\(93\)90258-z](https://doi.org/10.1016/0044-8486(93)90258-z)

Fuller, R. C. (1998). Fecundity estimates for rainbow darters, *Etheostoma caeruleum*, in southwestern Michigan. *Ohio Journal of Science* **98**, 2–5.

Fuller, R. C. (2003). Disentangling female mate choice and male competition in the rainbow darter, *Etheostoma caeruleum*. *Copeia* **2003**, 138–148. [https://doi.org/10.1643/0045-8511\(2003\)003\[0138:dfncam\]2.0.co;2](https://doi.org/10.1643/0045-8511(2003)003[0138:dfncam]2.0.co;2)

de Gaudemar, B. & Beall, E. (1998). Effects of overripening on spawning behaviour and reproductive success of Atlantic salmon females spawning in a controlled flow channel. *Journal of Fish Biology* **53**, 434–446. <https://doi.org/10.1006/jfbi.1998.0716>

Janetos, A. C. (1980). Strategies of female mate choice: a theoretical analysis. *Behavioral Ecology and Sociobiology* **7**, 107–112. <https://doi.org/10.1007/bf00299515>

Jennions, M. D. & Petrie, M. (1997). Variation in mate choice and mating preferences: a review of causes and consequences. *Biological Reviews* **72**, 283–327. <https://doi.org/10.1017/s0006323196005014>

Kjørsvik, E., Mangor-Jensen, A. & Holmefjord, I. (1990). Egg quality in fishes. *Advances in Marine Biology* **26**, 71–113. [https://doi.org/10.1016/S0065-2881\(08\)60199-6](https://doi.org/10.1016/S0065-2881(08)60199-6)

Kokko, H., Brooks, R., Jennions, M. D. & Morley, J. (2003). The evolution of mate choice and mating biases. *Proceedings of the Royal Society B* **270**, 653–664. <https://doi.org/10.1098/rspb.2002.2235>

McEvoy, L. A. (1984). Ovulatory rhythms and overripening of eggs in cultivated turbot, *Scophthalmus maximus* L. *Journal of Fish Biology* **24**, 437–448. <https://doi.org/10.1111/j.1095-8649.1984.tb04814.x>

Mendelson, T. C. (2003). Sexual isolation evolves faster than hybrid inviability in a diverse and sexually dimorphic genus of fish (Percidae: *Etheostoma*). *Evolution* **57**, 317–327. [https://doi.org/10.1554/0014-3820\(2003\)057\[0317:sie\]2.0.co;2](https://doi.org/10.1554/0014-3820(2003)057[0317:sie]2.0.co;2)

Moran, R. L., von Ende, C. N. & King, B. H. (2013). Mate choice copying in two species of darters (Percidae: *Etheostoma*). *Behaviour* **150**, 1255–1274. <https://doi.org/10.1163/1568539X-00003092>

Moran, R. L., Zhou, M., Catchen, J. M. & Fuller, R. C. (2017). Male and female contributions to behavioral isolation in darters as a function of genetic distance and color distance. *Evolution* **71**, 2428–2444. <https://doi.org/10.1111/evo.13321>

Page, L. M. (1983). *The Handbook of Darters*. Neptune City, NJ: TFH Publications.

Pyron, M. (1995). Mating patterns and a test for female mate choice in *Etheostoma spectabile* (Pisces, Percidae). *Behavioral Ecology and Sociobiology* **36**, 407–412. <https://doi.org/10.1007/bf00177336>

Real, L. (1990). Search theory and mate choice. I. Models of single-sex discrimination. *The American Naturalist* **136**, 376–405. <https://doi.org/10.1086/285103>

Reynolds, J. D. & Gross, M. R. (1990). Costs and benefits of female mate choice: is there a lek paradox? *The American Naturalist* **136**, 230–243. <https://doi.org/10.1086/285093>

Williams, T. H. & Mendelson, T. C. (2010). Behavioral isolation based on visual signals in a sympatric pair of darter species. *Ethology* **116**, 1038–1049. <https://doi.org/10.1111/j.1439-0310.2010.01816.x>

Williams, T. H., Gumm, J. M. & Mendelson, T. C. (2013). Sexual selection acting on a speciation trait in darters (Percidae: *Etheostoma*). *Behavioral Ecology* **24**, 1407–1414. <https://doi.org/10.1093/beheco/art080>

Winn, H. E. (1958). Observation on the reproductive habits of darters (Pisces—Percidae). *American Midland Naturalist* **59**, 190–212. <https://doi.org/10.2307/2422384>

Wong, B. & Candolin, U. (2005). How is female mate choice affected by male competition? *Biological Reviews* **80**, 559–571. <https://doi.org/10.1017/s1464793105006809>

Zhou, M. & Fuller, R. C. (2014). Reproductive isolation between two darter species is enhanced and asymmetric in sympatry. *Journal of Fish Biology* **84**, 1389–1400. <https://doi.org/10.1111/jfb.12364>

Zhou, M., Loew, E. R. & Fuller, R. C. (2015). Sexually asymmetric colour-based species discrimination in orangethroat darters. *Animal Behaviour* **106**, 171–179. <https://doi.org/10.1016/j.anbehav.2015.05.016>